

## Fundamental class - definition\*

MATTHIAS KRECK

### 1. THE INTEGRAL FUNDAMENTAL CLASS

For compact manifolds one can characterize the orientability by the existence of a certain homology class called the fundamental class. The background is the following theorem. Recall that if  $M$  is an  $n$ -dimensional topological manifold (possibly with boundary), then for each  $x$  in the interior of  $M$ , one has  $H_n(M, M - x; \mathbb{Z}) \cong \mathbb{Z}$ , [2, 22.1].

**Theorem 1.1.** *Let  $M$  be an  $n$ -dimensional compact topological manifold (possibly with boundary). If  $M$  is connected and orientable then for each  $x$  in the interior of  $M$ , the map induced by the inclusion*

$$H_n(M, \partial M; \mathbb{Z}) \rightarrow H_n(M, M - x; \mathbb{Z}),$$

*is an isomorphism. In particular,  $H_n(M, \partial M; \mathbb{Z}) \cong \mathbb{Z}$ . If  $M$  is connected and non-orientable then  $H_n(M, \partial M; \mathbb{Z})$  is zero.*

*Proof.* If  $M$  is closed then this is part of [1, VIII Corollary 3.4]; see also [2, 22.26]. If  $M$  has a boundary  $\partial M$ , then the inclusion  $\partial M = \partial M \times \{1\} \subset M$  extends to an embedding  $\partial M \times I \subset M$  of a collar, where  $I = [0, 1]$  [3, Proposition 3.42]. Let  $M_0 = \text{cl.}(M - \partial M \times I)$ , so that  $M = M_0 \cup_{\partial M} (\partial M \times I)$ . By excision

$$H_n(M, \partial M; \mathbb{Z}) \cong H_n(M, \partial M \times I; \mathbb{Z}) \cong H_n(M - \partial M \times \{1\}, \partial M \times [0, 1]; \mathbb{Z}).$$

Now apply [1, VIII Corollary 3.4] to the open manifold  $X := M - \partial M \times \{1\}$  and the closed subset  $M_0 \subset X$ .  $\square$

Theorem 1.1 implies that a connected compact manifold  $M$  is orientable if and only if

$$H_n(M, \partial M; \mathbb{Z}) \cong \mathbb{Z}.$$

A choice of a generator is then called a **fundamental class**  $[M, \partial M] \in H_n(M, \partial M; \mathbb{Z})$  for  $M$ . The fundamental class determines by the isomorphism above a continuous choice of local orientations and in turn the fundamental class is determined by a homological orientation of  $M$ . In other words a connected compact manifold together with the choice of a fundamental class  $[M, \partial M]$  is the same as an oriented manifold. If  $M$  is not connected, then  $M$  is orientable if and only if all components are orientable. If the components are oriented the fundamental classes of the components give the **fundamental class of  $M$**  under the isomorphism which decomposes the homology groups into the homology groups of the components. Thus for oriented manifolds again one has a fundamental class which corresponds to an orientation as in the connected case. The construction of the fundamental class of an oriented closed

---

\*Atlas page: [www.map.mpim-bonn.mpg.de/Fundamental\\_class](http://www.map.mpim-bonn.mpg.de/Fundamental_class)

manifold is done inductively over an atlas (similarly for manifolds with boundary). Namely one has the following generalization of Theorem 1:

**Theorem 1.2** ([2, 22.24]). *Let  $M$  be a connected oriented  $n$ -dimensional manifold. Then for each compact subset  $K \subset M$  there is a class  $[M]_K \in H_n(M, M - K)$  such that the following hold.*

- (1) *If  $K \subset K'$  is another compact subset, then  $[M]_K'$  maps to  $M_K$  under the map induced by the inclusion.*
- (2) *For each  $x \in M$  the class  $[M]_x$  is the local orientation of  $M$ .*
- (3) *The classes  $M_K$  are uniquely characterized by these properties.*

Using this one can use the Mayer-Vietoris sequence to ‘glue’ together the local orientations inductively over a finite oriented atlas together to construct  $M_K$ . The inductive construction is rather indirect. If one defines the homology of a space  $X$  as the bordism classes of certain stratified spaces  $\mathcal{S}$  together with a continuous map to  $X$ , e.g. stratifolds, then the the fundamental class is easy to obtain, it is a tautology. Then the fundamental class of a closed manifold is the bordism class represented by the identity map

$$id : M \rightarrow M.$$

For this see [4, Chapter 7, Section 1].

## 2. THE $\mathbb{Z}/2$ -FUNDAMENTAL CLASS

For all  $n$ -dimensional connected compact manifolds - even if they are not orientable - one has

$$H_n(M, \partial M; \mathbb{Z}/2) = \mathbb{Z}/2,$$

and one calls the non-trivial element the  $\mathbb{Z}/2$ -fundamental class [1, VIII Definition 4.1]. As for the integral fundamental class (if  $M$  is oriented) one gets from these classes the  **$\mathbb{Z}/2$ -fundamental class** of a non-connected compact manifold. Also one has a generalization of Theorem 1.2 to non-compact connected manifolds, i.e. for each compact subset  $K$  one has  $H_n(M, M - K; \mathbb{Z}/2) \cong \mathbb{Z}/2$  and for  $K \subset K'$  the map induced by the inclusion is an isomorphism

$$H_n(M, M - K'; \mathbb{Z}/2) \rightarrow H_n(N, M - K; \mathbb{Z}/2).$$

## REFERENCES

- [1] A. Dold, *Lectures on algebraic topology*, Springer-Verlag, 1995. [MR 1335915](#) [Zbl 0872.55001](#)
- [2] M. J. Greenberg and J. R. Harper, *Algebraic topology*, Benjamin/Cummings Publishing Co. Inc. Advanced Book Program, 1981. [MR 643101](#) [Zbl 0498.55001](#)
- [3] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002. [MR 1867354](#) [Zbl 1044.55001](#)
- [4] M. Kreck, *Differential algebraic topology*, Graduate Studies in Mathematics, 110, American Mathematical Society, 2010. [MR 2641092](#) [Zbl 05714474](#)

MATTHIAS KRECK  
 ENDENICHER ALLEE 60  
 D-53115 BONN, GERMANY

*E-mail address:* [kreck@math.uni-bonn.de](mailto:kreck@math.uni-bonn.de)

*Web address:* <http://him.uni-bonn.de/homepages/prof-dr-matthias-kreck/>